

USMT-06 - Mathematics-II Paper-VI : Set Theory and Laplace Transform

P. Pages : 2

Time : Three Hours



GUG/S/23/11613

Max. Marks : 60

- Notes : 1. Solve all **five** questions.
2. All questions carry equal marks.

UNIT – I

1. a) Prove that $(A \cup B)' = A' \cap B'$. 6
- b) Show that if $a, b \in \mathbb{R}$ then there is one & only one number x such that $x + a = b$. 6
This number x is given by $x = b + (-a)$
- OR**
- c) Show that $x \in \mathbb{R} \Rightarrow x^2 \geq 0$. 6
- d) Prove that Let $\langle A_n \rangle$ be a sequence of countable sets. Then $S = \bigcup_{n=1}^{\infty} A_n$ is countable. 6

UNIT – II

2. a) Prove that A fuzzy set F on the universal set R is convex if and only if $F[\lambda x_1 + (1-\lambda)x_2] \geq \min[F(x_1), F(x_2)] \forall x_1, x_2 \in R$ & all $\lambda \in [0, 1]$. 6
- b) Find $\tilde{A} \tilde{B}$, where 6

$$\tilde{A} = \frac{0.9}{1} + \frac{0.7}{3} + \frac{0.2}{4} + \frac{0.3}{6},$$

$$\tilde{B} = \frac{0.1}{2} + \frac{0.4}{3} + \frac{0.5}{4} + \frac{0.8}{5}$$
 are defined on the universe $U = \{1, 2, 3, 4, 5, 6\}$

OR

- c) Let $\tilde{A}, \tilde{B} \in P(U)$ Then for all $\alpha \in [0, 1]$ show that 6
- i) $\tilde{A} = \tilde{B}$ if and only if ${}^{\alpha}\tilde{A} = {}^{\alpha}\tilde{B}$
- ii) $\tilde{A} = \tilde{B}$ if and only if ${}^{\alpha+}\tilde{A} = {}^{\alpha+}\tilde{B}$
- d) Let $\tilde{A} \in P(U)$ then prove that ${}^{\alpha}(A') = [{}^{(1-\alpha)+}A]'$, $\forall \alpha \in [0, 1]$. 6

UNIT – III

3. a) Show that if $f(t)$ is piecewise continuous in every finite interval $0 \leq t \leq T$ for every $T > 0$ & of exponential order α for $t > T$ then its Laplace Transform $F(s)$ exists $\forall S > \alpha$. 6
- b) Find the Laplace Transform of function $\cosh at \cdot \sin at$. 6

OR

c) Find Laplace transform of 6

$$\cosh t \int_0^t e^t \cosh t dt$$

d) Obtain the Laplace Transform of the 6

$$x'' + 2x' - 3x, x(0) = 0, x'(0) = 1$$

UNIT – IV

4. a) Find the solution by Laplace Transform of $\frac{d^2x}{dt^2} + 9x = \cos 2t, x(0) = 1, x(\pi/2) = -1$. 6

b) Find the inverse Laplace Transform of $\frac{s^2 - 6}{s^3 + 4s^2 + 3s}$ 6

OR

c) Solve equation 6

$$\frac{d^2x}{dt^2} - 3\frac{dx}{dt} + 2x = 4t + e^{3t}$$

when $x(0) = 1$ and $x'(0) = -1$.

d) Find inverse Laplace Transform of $\frac{1}{(s-2)(s+2)^2}$ by convolution theorem. 6

5. Attempt **any six**.

a) Prove that 2
 $a + b = a + c \Rightarrow b = c \forall a, b, c \in \mathbb{R}$

b) State law of trichotomy. 2

c) Define intersection of two fuzzy set. 2

d) Define union of crisp set. 2

e) Find : $L[3t^2 - 2 \cdot e^t]$ 2

f) If $L[f(t)] = F(s)$ then prove that $L[e^{at} f(t)] = F(s-a)$. 2

g) Find Inverse Laplace transform of $\frac{s^2 - 3s + 4}{s^3}$. 2

h) State the convolution theorem. 2
